

Hashimoto's Encephalopathy: Back to Square One

Lord Brain and colleagues' classic description is cautiously entitled "Hashimoto's Disease and Encephalopathy." At the end of their discussion, the authors concluded that "antibody studies in future cases of unexplained encephalopathy should show whether we have described a syndrome or a coincidence."¹ At present, more than 250 additional adult cases with "Hashimoto's encephalopathy" (HE) have been reported in the literature, revealing a wide clinical spectrum, including movement disorders such as myoclonus (>20% of patients), ataxia, tremor, and opsoclonus.^{2,3} It has been suggested that this condition be referred to as steroid-responsive encephalopathy and associated autoimmune thyroiditis (SREAT), given that the role of anti-thyroid autoantibodies (ATAs) is far from clear.² However, at the same time, this new name implies steroid responsiveness, which is not always the case.^{2,4}

In an effort to solve the riddle that this condition entails, Mattozzi and colleagues recently conducted a study to determine whether pretreatment diagnostic criteria for HE predict steroid responsiveness.⁴ They included 24 patients (14 women) with suspected HE and only six of them completely responded to steroids.⁴ In addition, the authors reported that serum thyroid peroxidase (TPO) autoantibodies had a very limited specificity for this condition, because they were also detected in 8.1% and 8.2% of patients with possible autoimmune encephalitis and control subjects, respectively.⁴ Finally, only one patient with suspected HE had autoantibodies against the amino-terminal of alpha-enolase (NAE),⁴ a serum biomarker that had previously been reported to have high specificity for this condition.³ NAE autoantibodies were also detected in one control subject.⁴

In view of these findings, it seems we are back at square one. Considering the lack of steroid responsiveness in the majority of cases, the term SREAT seems to be a misnomer. Given ATAs are not specific for the disease, "Hashimoto's disease and encephalopathy" (rather than HE) seems an appropriate description. Most likely, Hashimoto's disease and encephalopathy is not a single disease entity but a clinical spectrum of as yet unexplained encephalopathies with different underlying pathomechanisms. ATAs probably only reflect a general susceptibility to autoimmunity, as Becker and colleagues suggested more than 5 decades ago.⁵

For the present, it is important to keep this in mind and search for disease-defining autoantibodies in any such patient. Particularly, although not exclusively, encephalitis with GABA_AR-

autoantibodies frequently associates with thyroid autoimmunity. Further studies to define homogenous subgroups will be needed to find specific biomarkers and to understand the underlying pathophysiological mechanisms.

Acknowledgment

G. D.-G. thanks the Consejo Nacional de Ciencia y Tecnología (Conacyt) for the support granted through a Full National Scholarship to pursue his MSc degree (CVU: 613905).

Author Roles

(1) Research Project: A. Conception, B. Organization, C. Execution; (2) Statistical Analysis: A. Design, B. Execution, C. Review and Critique; (3) Manuscript: A. Writing of the First Draft, B. Review and Critique.

G.D.-G.: 3A.

B.B.: 1A, 3B.

Disclosures

Ethical Compliance Statement: The authors confirm that the approval of an institutional review board and patient consent were not required for this work. We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this work is consistent with those guidelines.

Funding Sources and Conflicts of Interest: No specific funding was received for this work. The authors declare that there are no conflicts of interest relevant to this work.

Financial Disclosures for previous 12 months: The authors declare that there are no additional disclosures to report. ■

Guillermo Delgado-García MD,^{1,2}  Bettina Balint MD^{3,4}

¹Laboratorio Clínico de Enfermedades Neurodegenerativas, Instituto Nacional de Neurología y Neurocirugía, Mexico City, Mexico,

²Programa de Maestría y Doctorado en Ciencias Médicas, Odontológicas y de la Salud (PMDCMOS), Universidad Nacional

***Correspondence to:** Dr. Bettina Balint, Department of Neurology, University Hospital Heidelberg, Heidelberg, Germany; E-mail: bettina.balint@med.uni-heidelberg.de

Keywords: Hashimoto's encephalopathy, steroid-responsive encephalopathy associated with autoimmune thyroiditis, autoimmune encephalitis, myoclonus, Hashimoto's thyroiditis.

Mattozzi, S, Sabater, L, Escudero, D, et al. *Hashimoto encephalopathy in the 21st century. Neurology* 2020; 94: e217–e224.

Received 14 May 2020; revised 12 June 2020; accepted 14 June 2020.

Published online 16 July 2020 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mdc3.13012

Autónoma de México, Mexico City, Mexico, ³Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology, London, UK; ⁴Department of Neurology, University Hospital Heidelberg, Heidelberg, Germany

References

1. Brain L, Jellinek EH, Ball K. Hashimoto's disease and encephalopathy. *Lancet* 1966;2:512–514.
2. Laurent C, Capron J, Quillerou B, Thomas G, Alamowitch S, Fain O, Mekinian A. Steroid-responsive encephalopathy associated with autoimmune thyroiditis (SREAT): characteristics, treatment and outcome in 251 cases from the literature. *Autoimmun Rev* 2016;15:1129–1133.
3. Zhou JY, Xu B, Lopes J, Blamoun J, Li L. Hashimoto encephalopathy: literature review. *Acta Neurol Scand* 2017;135:285–290.
4. Mattozzi S, Sabater L, Escudero D, et al. Hashimoto encephalopathy in the 21st century. *Neurology* 2020;94:e217–e224.
5. Becker KL, Titus JL, Woolner LB, et al. Significance of morphologic thyroiditis. *Ann Intern Med* 1965;62:1134–1138.